

# MIXED TORIC RESIDUES AND CALABI-YAU COMPLETE INTERSECTIONS

VICTOR V. BATYREV AND EVGENY N. MATEROV

ABSTRACT. Using the Cayley trick, we define the notions of mixed toric residues and mixed Hessians associated with  $r$  Laurent polynomials  $f_1, \dots, f_r$ . We conjecture that the values of mixed toric residues on the mixed Hessians are determined by mixed volumes of the Newton polytopes of  $f_1, \dots, f_r$ . Using mixed toric residues, we generalize our Toric Residue Mirror Conjecture to the case of Calabi-Yau complete intersections in Gorenstein toric Fano varieties obtained from nef-partitions of reflexive polytopes.

## CONTENTS

|                                                           |    |
|-----------------------------------------------------------|----|
| 1. Introduction                                           | 2  |
| 2. Toric residues                                         | 3  |
| 3. The Cayley trick and mixed toric residues              | 7  |
| 4. Toric Residue Mirror Conjecture                        | 11 |
| 5. Complete intersections in weighted projective spaces   | 14 |
| 6. Complete intersections in product of projective spaces | 17 |
| 7. Computation of Yukawa $(d - r)$ -point functions       | 20 |
| References                                                | 27 |

---

*Key words and phrases.* residues, toric varieties, intersection numbers, mirror symmetry.  
 2000 *Mathematics Subject Classification.* Primary 14M25.

## 1. INTRODUCTION

This paper is the continuation of our previous work [BM1] where we proposed a toric mirror symmetry test using toric residues. The idea of this test has appeared in the paper of Morrison-Plesser [MP] who observed that the coefficients of some power series expansions of unnormalized Yukawa couplings for mirrors of Calabi-Yau hypersurfaces in toric varieties  $\mathbb{P}$  can be interpreted as generating functions for intersection numbers of divisors on some sequences of toric varieties  $\mathbb{P}_\beta$  parameterized by lattice points  $\beta$  in the Mori cone  $K_{\text{eff}}(\mathbb{P})$  of  $\mathbb{P}$ . Due to results of Mavlyutov [Mav2], it is known that the unnormalized Yukawa couplings can be computed using the toric residues introduced by Cox [Cox]. In our paper [BM1], we formulated a general mathematical conjecture, so called *Toric Residue Mirror Conjecture*, which describes some power series expansions of the toric residues in terms of intersection numbers of divisors on a sequence of simplicial toric varieties  $\mathbb{P}_\beta$  (we call them Morrison-Plesser moduli spaces). This conjecture includes all examples of mirror symmetry for Calabi-Yau hypersurfaces in Gorenstein toric varieties associated with reflexive polytopes. Since the toric mirror symmetry construction exists also for Calabi-Yau complete intersection in Gorenstein toric Fano varieties [Bo, BB1], it is natural to try to extend our conjecture to this more general situation.

The case of Calabi-Yau complete intersections of  $r$  hypersurfaces

$$f_1(t) = \cdots = f_r(t) = 0, \quad r > 1,$$

defined by Laurent polynomials  $f_1(t), \dots, f_r(t) \in \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$  in  $d$ -dimensional toric varieties  $\mathbb{P}$  was not considered by Morrison and Plesser in [MP]. We remark that in this case one *does not* get a connection to the “quantum cohomology ring” [Bat] as in the hypersurface case. This difference is explained by the consideration of a non-reflexive  $(d+r-1)$ -dimensional polytope  $\tilde{\Delta}$ , so called Cayley polytope, and its secondary polytope  $\text{Sec}(\tilde{\Delta})$ . The Cayley polytope  $\tilde{\Delta}$  appears from the Cayley trick which introduces  $r$  additional  $r$  variables  $t_{d+1}, \dots, t_{d+r}$  and a new polynomial  $F(t) := \sum_{j=1}^r t_{d+j} f_j(t)$ . We consider the usual toric residue  $\text{Res}_F$  associated with  $F$  and define the  $k$ -mixed toric residue  $\text{Res}_F^k$  corresponding to a positive integral solution  $k = (k_1, \dots, k_r)$  of the equation  $k_1 + \cdots + k_r = d+r$  as a  $k$ -th homogeneous component of  $\text{Res}_F$ . We expect that the  $k$ -mixed toric residues are similar to the usual toric residues. In particular, we introduce the notion of the  $k$ -mixed Hessian  $H_F^k$  of Laurent polynomials  $f_1, \dots, f_r$  and conjecture that the value of  $\text{Res}_F^k$  on  $H_F^k$  is exactly the mixed volume

$$V(\underbrace{\Delta_1, \dots, \Delta_1}_{k_1-1}, \dots, \underbrace{\Delta_r, \dots, \Delta_r}_{k_r-1}),$$

where  $\Delta_1, \dots, \Delta_r$  are Newton polytopes of  $f_1, \dots, f_r$ .

Our generalization of the Toric Residue Mirror Conjecture for Calabi-Yau complete intersections uses the notions of the nef-partition  $\Delta = \Delta_1 + \dots + \Delta_r$  of  $d$ -dimensional reflexive polytope  $\Delta$  [Bo]. In this situation, one obtains a dual nef-partition  $\nabla = \nabla_1 + \dots + \nabla_r$  and two more reflexive polytopes:

$$\nabla^* = \text{conv}\{\Delta_1, \dots, \Delta_r\}, \quad \Delta^* = \text{conv}\{\nabla_1, \dots, \nabla_r\}.$$

It is important that special coherent triangulations of  $\nabla^*$  define coherent triangulations of the Cayley polytope  $\tilde{\Delta} := \Delta_1 * \dots * \Delta_r$ . Therefore the choice of such a triangulation  $\mathcal{T}$  of  $\nabla^*$  determines a vertex  $v_{\mathcal{T}}$  of the secondary polytope  $\text{Sec}(\tilde{\Delta})$  and a partial projective simplicial crepant desingularization  $\mathbb{P} := \mathbb{P}_{\Sigma(\mathcal{T})}$  of the Gorenstein toric variety  $\mathbb{P}_{\nabla^*}$ . So one obtains a sequence of simplicial toric varieties  $\mathbb{P}_{\beta}$  associated with the lattice points  $\beta$  in the Mori cone  $K_{\text{eff}}(\mathbb{P})$  of  $\mathbb{P}$ . We conjecture that the generating function of intersection numbers

$$I_P(a) = \sum_{\beta \in K_{\text{eff}}(\mathbb{P})} I(P, \beta) a^{\beta}$$

coincides with the power series expansion of the  $k$ -mixed toric residue

$$R_P(a) = \text{Res}_F^k(P(a, t))$$

at the vertex  $v_{\mathcal{T}} \in \text{Sec}(\tilde{\Delta})$ . The precise formulation of this conjecture is given in Section 4.

In Sections 5, 6 we check our conjecture for nef-partitions corresponding to Calabi-Yau complete intersections in weighted projective spaces  $\mathbb{P}(w_1, \dots, w_n)$  and in product of projective spaces  $\mathbb{P}^{d_1} \times \dots \times \mathbb{P}^{d_p}$ . The final section is devoted to applications of the Toric Residue Mirror Conjecture to the computation of Yukawa couplings for Calabi-Yau complete intersections.

*Acknowledgments.* This work was supported by DFG, Forschungsschwerpunkt ‘‘Globale Methoden in der komplexen Geometrie’’. E. Materov was supported in part by RFBR Grant 00-15-96140. We are also appreciate to hospitality and support of the Isaac Newton Institute in Cambridge.

## 2. TORIC RESIDUES

In this section we remind necessary well-known facts about toric residues (see [Cox, CDS, BM1]).

Let  $\tilde{M}$  and  $\tilde{N} = \text{Hom}(\tilde{M}, \mathbb{Z})$  be two free abelian groups of rank  $\tilde{d}$  dual to each other. We denote by

$$\langle *, * \rangle : \tilde{M} \times \tilde{N} \rightarrow \mathbb{Z}$$

the natural bilinear pairing, and by  $\tilde{M}_{\mathbb{R}}$  (resp.  $\tilde{N}_{\mathbb{R}}$ ) the real scalar extension of  $\tilde{M}$  (resp.  $\tilde{N}$ ).

**Definition 2.1** ([BB2]). A  $\tilde{d}$ -dimensional rational polyhedral cone  $C$  ( $\tilde{d} > 0$ ) in  $\widetilde{M}_{\mathbb{R}}$  is called *Gorenstein* if it is strongly convex (i.e.,  $C + (-C) = \{0\}$ ), there exists an element  $n_C \in \tilde{N}$  such that  $\langle x, n_C \rangle > 0$  for any nonzero  $x \in C$ , and all vertices of the  $(\tilde{d} - 1)$ -dimensional convex polytope

$$\Delta(C) = \{x \in C : \langle x, n_C \rangle = 1\}$$

belong to  $\widetilde{M}$ . The polytope  $\Delta(C)$  is called the *supporting polytope* of  $C$ . For any  $m \in C \cap \widetilde{M}$ , we define the *degree* of  $m$  as

$$\deg m = \langle m, n_C \rangle.$$

**Definition 2.2.** Let  $\tilde{\Delta} = \Delta(C)$  be the supporting polytope for a Gorenstein cone  $C \subset \widetilde{M}_{\mathbb{R}}$ . We denote by  $S_{\tilde{\Delta}}$  the semigroup  $\mathbb{C}$ -algebra of the monoid of lattice points  $C \cap \widetilde{M}$ . In order to transform the additive semigroup operation in  $C \cap \widetilde{M}$  into a multiplicative form in  $S_{\tilde{\Delta}}$ , we write  $t^m$  for the element in  $S_{\tilde{\Delta}}$  corresponding to  $m \in C$ . One can consider  $S_{\tilde{\Delta}}$  as a graded  $\mathbb{C}$ -algebra:

$$S_{\tilde{\Delta}} = \bigoplus_{l=0}^{\infty} S_{\tilde{\Delta}}^{(l)},$$

where the  $l$ -th homogeneous component  $S_{\tilde{\Delta}}^{(l)}$  has a  $\mathbb{C}$ -basis consisting of all  $t^m$  such that  $m \in C \cap \widetilde{M}$  and  $\deg m = l$ . We define also the homogeneous ideal

$$I_{\tilde{\Delta}} = \bigoplus_{l=0}^{\infty} I_{\tilde{\Delta}}^{(l)}$$

in  $S_{\tilde{\Delta}}$  whose  $\mathbb{C}$ -basis consists of all  $t^m$  such that  $m$  is a lattice point in the interior of  $C$ .

**Definition 2.3.** An element

$$g := \sum_{m \in \tilde{\Delta} \cap \widetilde{M}} a_m t^m \in S_{\tilde{\Delta}}^{(1)}, \quad a_m \in \mathbb{C}$$

is called  $\tilde{\Delta}$ -regular if for some  $\mathbb{Z}$ -basis  $n_1, \dots, n_{\tilde{d}}$  of  $\tilde{N}$  the elements

$$g_i := \sum_{m \in \tilde{\Delta} \cap \widetilde{M}} a_m \langle m, n_i \rangle t^m, \quad i = 1, \dots, \tilde{d}$$

form a regular sequence in  $S_{\tilde{\Delta}}$ . We define the matrix  $G := (g_{ij})_{1 \leq i, j \leq \tilde{d}}$ , where

$$g_{ij} := \sum_{m \in \tilde{\Delta} \cap \widetilde{M}} a_m \langle m, n_i \rangle \langle m, n_j \rangle t^m, \quad i, j = 1, \dots, \tilde{d}.$$

The element

$$H_g := \det G$$

is called *Hessian of  $g$* .

*Remark 2.4.* a) The definition of  $\tilde{\Delta}$ -regularity does not depend on the choice of  $\mathbb{Z}$ -basis  $n_1, \dots, n_{\tilde{d}}$  of  $\tilde{N}$ . It gives the same non-degeneracy condition as the one proposed by Mavlyutov in [Mav1]. In many applications the lattice vector  $n_C$  will be included in  $\{n_1, \dots, n_{\tilde{d}}\}$ .

b) If  $\tilde{\Delta} \cap \tilde{M} = \{m_1, \dots, m_{\mu}\}$ , then by [CDS, Proposition 1.2], one has

$$H_g = \sum_{1 \leq i_1 < \dots < i_{\tilde{d}} \leq \mu} (\det(m_{i_1}, \dots, m_{i_{\tilde{d}}}))^2 t^{m_{i_1} + \dots + m_{i_{\tilde{d}}}}.$$

In particular,  $H_g$  is independent on the choice of the  $\mathbb{Z}$ -basis  $n_1, \dots, n_{\tilde{d}}$  and  $H_g \in I_{\tilde{\Delta}}^{(\tilde{d})}$ .

c) The graded  $\mathbb{C}$ -algebra  $S_{\tilde{\Delta}}$  is Cohen-Macaulay and  $I_{\tilde{\Delta}}$  is its dualizing module. If  $g$  is  $\tilde{\Delta}$ -regular in  $S_{\tilde{\Delta}}$ , then

$$S_g := S_{\tilde{\Delta}} / \langle g_1, \dots, g_{\tilde{d}} \rangle S_{\tilde{\Delta}},$$

is a graded finite-dimensional ring and

$$I_g := I_{\tilde{\Delta}} / \langle g_1, \dots, g_{\tilde{d}} \rangle I_{\tilde{\Delta}}$$

is a graded  $S_g$ -module together with a non-degenerate pairing

$$S_g^{(l)} \times I_g^{(\tilde{d}-l)} \rightarrow I_g^{(\tilde{d})} \simeq \mathbb{C}, \quad l = 0, \dots, \tilde{d} - 1.$$

induced by the  $S_g$ -module structure.

**Definition 2.5.** By the *toric residue* corresponding to a  $\tilde{\Delta}$ -regular element  $g \in S_{\tilde{\Delta}}^{(1)}$  we mean the  $\mathbb{C}$ -linear mapping

$$\text{Res}_g : I_{\tilde{\Delta}}^{(\tilde{d})} \rightarrow \mathbb{C}$$

which is uniquely determined by two conditions:

- (i)  $\text{Res}_g(h) = 0$  for any  $h \in \langle g_1, \dots, g_{\tilde{d}} \rangle I_{\tilde{\Delta}}$ ;
- (ii)  $\text{Res}_g(H_g) = \text{Vol}(\tilde{\Delta})$ , where  $\text{Vol}(\tilde{\Delta})$  denotes the volume of the  $(\tilde{d} - 1)$ -dimensional polytope  $\tilde{\Delta}$  multiplied by  $(\tilde{d} - 1)!$ .

Let  $\mathbb{P}_{\tilde{\Delta}} := \text{Proj } S_{\tilde{\Delta}}$  be the  $(\tilde{d} - 1)$ -dimensional toric variety associated with the polytope  $\tilde{\Delta}$  and  $\mathcal{O}_{\mathbb{P}_{\tilde{\Delta}}}(1)$  the corresponding ample sheaf on  $\mathbb{P}_{\tilde{\Delta}}$ . Then one has the canonical isomorphisms of graded rings

$$S_{\tilde{\Delta}} \cong \bigoplus_{l \geq 0} H^0(\mathbb{P}_{\tilde{\Delta}}, \mathcal{O}_{\mathbb{P}_{\tilde{\Delta}}}(l))$$

and graded modules

$$I_{\tilde{\Delta}} \cong \bigoplus_{l \geq 0} H^0(\mathbb{P}_{\tilde{\Delta}}, \omega_{\mathbb{P}_{\tilde{\Delta}}}(l)),$$

where  $\omega_{\mathbb{P}_{\tilde{\Delta}}}$  is the dualizing sheaf on  $\mathbb{P}_{\tilde{\Delta}}$ . In particular, we obtain a canonical isomorphism

$$I_{\tilde{\Delta}}^{(\tilde{d})} \cong H^0(\mathbb{P}_{\tilde{\Delta}}, \omega_{\mathbb{P}_{\tilde{\Delta}}}(\tilde{d})).$$

The following statement is a simple reformulation of Theorem 2.9(i) in [BM1]:

**Proposition 2.6.** *Let  $n_1, \dots, n_{\tilde{d}}$  be a  $\mathbb{Z}$ -basis of  $\tilde{N}$  such that  $n_1 = n_C$ . Denote by  $m_1, \dots, m_{\tilde{d}}$  the dual  $\mathbb{Z}$ -basis of  $\tilde{M}$ . For any elements  $h \in I_{\tilde{\Delta}}^{(\tilde{d})}$  and  $g \in S_{\tilde{\Delta}}^{(1)}$ , we define a rational differential  $(\tilde{d} - 1)$ -form on  $\mathbb{P}_{\tilde{\Delta}}$ :*

$$\Omega(h, g) := \frac{h}{g_1 \cdots g_{\tilde{d}}} \frac{dt^{m_2}}{t^{m_2}} \wedge \cdots \wedge \frac{dt^{m_{\tilde{d}}}}{t^{m_{\tilde{d}}}}.$$

If  $g$  is  $\tilde{\Delta}$ -regular, then

$$\text{Res}_g(h) = \sum_{\xi \in V_g} \text{res}_{\xi}(\Omega(h, g)),$$

where  $V_g = \{\xi \in \mathbb{P}_{\tilde{\Delta}} : g_2(\xi) = \cdots = g_{\tilde{d}}(\xi) = 0\}$  is the set of common zeros of  $g_2, \dots, g_{\tilde{d}}$  and  $\text{res}_{\xi}(\Omega(h, g))$  is the local Grothendieck residue of the form  $\Omega(h, g)$  at the point  $\xi \in V_g$ .

In particular, if all the common roots of  $g_2, \dots, g_{\tilde{d}}$  are simple and contained in the open dense  $(\tilde{d} - 1)$ -dimensional torus  $\mathbb{T} \subset \mathbb{P}_{\tilde{\Delta}}$ , then

$$\text{Res}_g(h) = \sum_{\xi \in V_g} \frac{p(\xi)}{g_1(\xi) H_g^1(\xi)},$$

where  $H_g^1$  is the determinant of the matrix  $G^1 := (g_{ij})_{2 \leq i, j \leq \tilde{d}}$ .

**Definition 2.7** ([BB1]). A Gorenstein cone  $C$  is called *reflexive* if the dual cone

$$\check{C} = \{y \in \tilde{N}_{\mathbb{R}} : \langle x, y \rangle \geq 0 \quad \forall x \in C\}$$

is also Gorenstein, i.e., there exists  $m_{\check{C}} \in \tilde{M}$  such that  $\langle m_{\check{C}}, y \rangle > 0$  for all  $y \in \check{C} \setminus \{0\}$ , and all vertices of the supporting polytope

$$\Delta(\check{C}) = \{y \in \check{C} : \langle m_{\check{C}}, y \rangle = 1\}$$

belong to  $\tilde{N}$ . We will call the integer  $r = \langle m_{\check{C}}, n_C \rangle$  the *index* of  $C$  (or  $\check{C}$ ). A  $(\tilde{d} - 1)$ -dimensional lattice polytope  $\tilde{\Delta}$  is called *reflexive* if it is a supporting polytope of some  $\tilde{d}$ -dimensional reflexive Gorenstein cone  $C$  of index 1. Moreover, the supporting polytope  $\tilde{\Delta}^*$  of the dual cone  $\check{C}$  is also reflexive polytope which is called *dual* (or *polar*) to  $\tilde{\Delta}$ .

If  $C$  is a reflexive Gorenstein cone of index  $r$ , then  $I_{\tilde{\Delta}}$  is a principal ideal generated by the element  $t^{m_{\tilde{c}}}$  of degree  $r$ . So one obtains the canonical isomorphism  $I_{\tilde{\Delta}}^{(l)} \cong S_{\tilde{\Delta}}^{(l-r)}$ . In particular, there exists the toric residue mapping

$$\text{Res}_g : S_{\tilde{\Delta}}^{(\tilde{d}-r)} \rightarrow \mathbb{C}$$

which is uniquely determined by the conditions:

- (i)  $\text{Res}_g(h) = 0$  for any  $h \in \langle g_1, \dots, g_{\tilde{d}} \rangle S_{\tilde{\Delta}}$ ;
- (ii)  $\text{Res}_g(H'_g) = \text{Vol}(\tilde{\Delta})$ , where  $H_g = t^{m_{\tilde{c}}} H'_g$ .

### 3. THE CAYLEY TRICK AND MIXED TORIC RESIDUES

Let  $M$  be a free abelian group of rank  $d$ ,  $M_{\mathbb{R}} := M \otimes \mathbb{R}$ , and  $\Delta \subset M_{\mathbb{R}}$  a convex  $d$ -dimensional polytope with vertices in  $M$ . We assume that there exist  $r$  convex polytopes  $\Delta_1, \dots, \Delta_r$  with vertices in  $M$  such that  $\Delta$  can be written as the Minkowski sum  $\Delta = \Delta_1 + \dots + \Delta_r$  (here we do not require that all polytopes  $\Delta_1, \dots, \Delta_r$  have maximal dimension  $d$ ).

**Definition 3.1.** We set  $\tilde{M} := M \oplus \mathbb{Z}^r$ ,  $\tilde{d} := d + r$  and define the  $\tilde{d}$ -dimensional Gorenstein cone  $C = C(\Delta_1, \dots, \Delta_r)$  in  $\tilde{M}_{\mathbb{R}} := M_{\mathbb{R}} \oplus \mathbb{R}^r$  as follows

$$C := \{(\lambda_1 x_1 + \dots + \lambda_r x_r, \lambda_1, \dots, \lambda_r) \in \tilde{M}_{\mathbb{R}} : \lambda_i \geq 0, x_i \in \Delta_i, i = 1, \dots, r\}.$$

The  $(d + r - 1)$ -dimensional polytope  $\Delta_1 * \dots * \Delta_r$  defined as the intersection of the cone  $C$  with the affine hyperplane  $\sum_{i=1}^r \lambda_i = 1$

$$\Delta_1 * \dots * \Delta_r := \{(\lambda_1 x_1 + \dots + \lambda_r x_r, \lambda_1, \dots, \lambda_r) : \lambda_i \geq 0, \sum_{i=1}^r \lambda_i = 1, x_i \in \Delta_i\},$$

will be called the *Cayley polytope associated with the Minkowski sum decomposition*  $\Delta = \Delta_1 + \dots + \Delta_r$ . It is clear that all vertices of  $\Delta_1 * \dots * \Delta_r$  are contained in  $\tilde{M}$  and

$$\Delta_1 * \dots * \Delta_r = \text{conv}((\Delta_1 \times \{b_1\}) \cup \dots \cup (\Delta_r \times \{b_r\})),$$

where  $\{b_1, \dots, b_r\}$  is the standard basis of  $\mathbb{Z}^r$ . For fixed polytopes  $\Delta_1, \dots, \Delta_r$  we denote  $\Delta_1 * \dots * \Delta_r$  simply by  $\tilde{\Delta}$ .

**Definition 3.2.** Define  $S_{\tilde{\Delta}} := \mathbb{C}[C \cap \tilde{M}]$  to be the semigroup algebra of the monoid  $C \cap \tilde{M}$  over the complex numbers. The algebra  $S_{\tilde{\Delta}}$  has a natural  $\mathbb{Z}_{\geq 0}^r$ -grading defined by the last  $r$  coordinates of lattice points in  $\tilde{M}$ . By choosing an isomorphism  $M \cong \mathbb{Z}^d$ , we can identify  $S_{\tilde{\Delta}}$  with a  $\mathbb{Z}_{\geq 0}^r$ -graded monomial subalgebra in

$$\mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}, t_{d+1}, \dots, t_{d+r}],$$

where the  $\mathbb{Z}_{\geq 0}^r$ -grading is considered with respect to the last  $r$  variables  $t_{d+1}, \dots, t_{d+r}$ . We denote by  $I_{\tilde{\Delta}}$  the  $\mathbb{Z}_{\geq 0}^r$ -graded monomial ideal in  $S_{\tilde{\Delta}}$  generated by all lattice points in the interior of  $C$ . For any  $k = (k_1, \dots, k_r) \in \mathbb{Z}_{\geq 0}^r$ , we denote by  $S_{\tilde{\Delta}}^k$  (resp. by  $I_{\tilde{\Delta}}^k$ ) the  $k$ -homogeneous component of  $S_{\tilde{\Delta}}$  (resp. of  $I_{\tilde{\Delta}}$ ). We will use also the total  $\mathbb{Z}_{\geq 0}$ -grading on  $S_{\tilde{\Delta}}$  and  $I_{\tilde{\Delta}}$ . For any nonnegative integer  $l$ , we denote the corresponding  $l$ -homogeneous components of  $S_{\tilde{\Delta}}$  and  $I_{\tilde{\Delta}}$  by  $S_{\tilde{\Delta}}^{(l)}$  and  $I_{\tilde{\Delta}}^{(l)}$  respectively. So one has:

$$S_{\tilde{\Delta}}^{(l)} = \bigoplus_{|k|=l} S_{\tilde{\Delta}}^k, \quad I_{\tilde{\Delta}}^{(l)} = \bigoplus_{|k|=l} I_{\tilde{\Delta}}^k,$$

where  $|k| := k_1 + \dots + k_r$ .

Let  $f_1(t), \dots, f_r(t)$  be Laurent polynomials in  $\mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$  such that  $\Delta_i$  is the Newton polytope of  $f_i$  ( $1 \leq i \leq r$ ). We set

$$F(t) := t_{d+1}f_1(t) + \dots + t_{d+r}f_r(t).$$

It is easy to see that  $\tilde{\Delta} = \Delta_1 * \dots * \Delta_r$  is the Newton polytope of  $F$ . Moreover, using the decomposition

$$S_{\tilde{\Delta}}^{(1)} = \bigoplus_{|k|=1} S_{\tilde{\Delta}}^k = \bigoplus_{i=1}^r S_{\tilde{\Delta}}^{b_i},$$

we see that every Laurent polynomial  $G$  in  $\mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}, t_{d+1}, \dots, t_{d+r}]$  with the Newton polytope  $\tilde{\Delta}$  can be obtained from the sequence of arbitrary Laurent polynomials  $g_1, \dots, g_r \in \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$  by the formula  $G = t_{d+1}g_1 + \dots + t_{d+r}g_r$ , where  $\Delta_i$  is the Newton polytope of  $g_i$  ( $1 \leq i \leq r$ ). The above correspondence  $\{f_1, \dots, f_r\} \mapsto F$  is usually called the *Cayley trick*. We call  $F = t_{d+1}f_1 + \dots + t_{d+r}f_r$  the *Cayley polynomial associated with  $f_1, \dots, f_r$* .

**Definition 3.3.** Let  $\Delta_1, \dots, \Delta_r \subset M_{\mathbb{R}}$  be convex polytopes with vertices in  $M$  such  $\Delta = \Delta_1 + \dots + \Delta_r$  has dimension  $d$ . We say that  $r$  Laurent polynomials

$$f_i(t) = \sum_{m \in \Delta_i \cap M} a_m^{(i)} t^m, \quad i = 1, \dots, r$$

form a  $\tilde{\Delta}$ -regular sequence if the corresponding Cayley polynomial  $F$  is  $\tilde{\Delta}$ -regular, i.e., the polynomials

$$F_i := t_i \partial / \partial t_i F, \quad i = 1, \dots, d+r$$

form a regular sequence in  $S_{\tilde{\Delta}}$ .

**Definition 3.4.** Let  $f_1(t), \dots, f_r(t)$  be Laurent polynomials with Newton polytopes  $\Delta_1, \dots, \Delta_r$  as above,  $F = t_{d+1}f_1 + \dots + t_{d+r}f_r$  the corresponding Cayley polynomial, and

$$H_F := \det \left( t_i \frac{\partial F_j}{\partial t_i} \right)_{1 \leq i, j \leq d+r} = \det \left( \left( t_i \frac{\partial}{\partial t_i} \right) \left( t_j \frac{\partial}{\partial t_j} \right) F \right)_{1 \leq i, j \leq d+r} \in I_{\tilde{\Delta}}^{(d+r)} \subset S_{\tilde{\Delta}}^{(d+r)}$$



the Hessian of  $F$ . For any  $k = (k_1, \dots, k_r)$  with  $|k| = d+r$  we define  $H_F^k \in I_{\tilde{\Delta}}^k$  to be the  $k$ -homogeneous component of  $H_F$ . The polynomial  $H_F^k$  will be called *k-mixed Hessian of  $f_1, \dots, f_r$* .

*Remark 3.5.* Since the last  $r$  rows of the matrix

$$\left( \left( t_i \frac{\partial}{\partial t_i} \right) \left( t_j \frac{\partial}{\partial t_j} \right) F \right)_{1 \leq i, j \leq d+r}$$

are divisible respectively by  $t_{d+1}, \dots, t_{d+r}$ , the Hessian  $H_F$  is divisible by the monomial  $t_{d+1} \cdots t_{d+r}$ . Therefore  $H_F^k = 0$  if one of the coordinates  $k_i$  of  $k = (k_1, \dots, k_r)$  is zero. In particular, one has

$$H_F = \sum_{\substack{k \in \mathbb{Z}_{>0}^r \\ |k|=d+r}} H_F^k.$$

Let  $k = (k_1, \dots, k_r) \in \mathbb{Z}_{>0}^r$  be a solution of the linear Diophantine equation

$$|k| = k_1 + \cdots + k_r = d + r.$$

For any  $r$  subsets  $S_i \subset \Delta_i \cap M$  such that  $|S_i| = k_i$  ( $1 \leq i \leq r$ ) we define the nonnegative integer  $\nu(S_1, \dots, S_r)$  as follows: choose an element  $s_i$  in each  $S_i$  ( $1 \leq i \leq r$ ), define  $S$  to be the  $d \times d$ -matrix whose rows are all possible nonzero vectors  $s - s_i$  where  $s \in S_i$ ,  $1 \leq i \leq r$ , and set  $\nu(S_1, \dots, S_r) := (\det S)^2$ . It is easy to see that up to sign  $\det S$  does not depend on the choice of elements  $s_i \in S_i$  and therefore  $\nu(S_1, \dots, S_r)$  is well defined.

**Proposition 3.6.** *Let  $k = (k_1, \dots, k_r) \in \mathbb{Z}_{>0}^r$  be a positive integral solution of the linear Diophantine equation*

$$|k| = k_1 + \cdots + k_r = d + r.$$

*Then the mixed Hessian can be computed by the following formula*

$$H_F^k = t_{d+1}^{k_1} \cdots t_{d+r}^{k_r} \sum_{(S_1, \dots, S_r)} \nu(S_1, \dots, S_r) \prod_{i=1}^r \prod_{s_i \in S_i} a_{s_i}^{(i)} t^{s_i},$$

where the sum runs over all  $r$ -tuples  $(S_1, \dots, S_r)$  of subsets  $S_i \subset \Delta_i \cap M$  such that  $|S_i| = k_i$  ( $1 \leq i \leq r$ ).

*Proof.* The formula for  $H_F^k$  follows immediately from the formula in 2.4(b) applied to the Cayley polytope  $\tilde{\Delta}$ .  $\square$

**Definition 3.7.** Let  $k = (k_1, \dots, k_r) \in \mathbb{Z}_{>0}^r$  be a positive integral solution of the equation

$$|k| = k_1 + \cdots + k_r = d + r.$$

Consider the toric residue

$$\text{Res}_F : I_{\tilde{\Delta}}^{(d+r)} \rightarrow \mathbb{C}$$

defined as a  $\mathbb{C}$ -linear map which vanishes on  $\langle F_1, \dots, F_{d+r} \rangle I_{\tilde{\Delta}}$  and sends  $H_F$  to  $\text{Vol}(\tilde{\Delta}) = \text{Vol}(\Delta_1 * \dots * \Delta_r)$ . The restriction  $\text{Res}_F^k$  of  $\text{Res}_F$  to the  $k$ -th homogeneous component  $I_{\tilde{\Delta}}^k$ :

$$\text{Res}_F^k : I_{\tilde{\Delta}}^k \rightarrow \mathbb{C}$$

will be called the *k-mixed toric residue associated with  $f_1, \dots, f_r$* .

Since  $H_F^k$  is an element of  $I_{\tilde{\Delta}}^k$ , it is natural to ask about the value of  $\text{Res}_F^k(H_F^k)$ .

**Conjecture 3.8.** *Let  $k = (k_1, \dots, k_r) \in \mathbb{Z}_{>0}^r$  be a positive integral solution of  $k_1 + \dots + k_r = d + r$ . We set  $\bar{k} = (\bar{k}_1, \dots, \bar{k}_r) := (k_1 - 1, \dots, k_r - 1)$ . Then*

$$\text{Res}_F^k(H_F^k) = V(\underbrace{\Delta_1, \dots, \Delta_1}_{\bar{k}_1}, \dots, \underbrace{\Delta_r, \dots, \Delta_r}_{\bar{k}_r}),$$

where  $V(\Theta_1, \dots, \Theta_d)$  denotes the mixed volume of convex polytopes  $\Theta_1, \dots, \Theta_d$  multiplied by  $(d + r - 1)!$ .

Our conjecture agrees with a result of Danilov and Khovanskii:

**Proposition 3.9.** [DK, §6] *The normalized volume of the Cayley polytope  $\tilde{\Delta} = \Delta_1 * \dots * \Delta_r$  can be computed by the following formula:*

$$\text{Vol}(\Delta_1 * \dots * \Delta_r) = \sum_{|\bar{k}|=d} V(\underbrace{\Delta_1, \dots, \Delta_1}_{\bar{k}_1}, \dots, \underbrace{\Delta_r, \dots, \Delta_r}_{\bar{k}_r}).$$

*Remark 3.10.* Let  $r = d$  and  $k = (d + 1, 1, \dots, 1)$ . It follows from 3.6 and 2.4(b) that

$$H_F^k = H_{t_{d+1}f_1}(t_{d+2}f_2) \cdots (t_{2d}f_d),$$

where

$$H_{t_{d+1}f_1} = \det \left( \left( t_i \frac{\partial}{\partial t_i} \right) \left( t_j \frac{\partial}{\partial t_j} \right) t_{d+1}f_1 \right)_{1 \leq i, j \leq d+1}.$$

On the other hand, we have

$$V(\underbrace{\Delta_1, \dots, \Delta_1}_{\bar{k}_1}, \dots, \underbrace{\Delta_r, \dots, \Delta_r}_{\bar{k}_r}) = V(\underbrace{\Delta_1, \dots, \Delta_1}_d) = \text{Vol}(\Delta_1).$$

Therefore, Conjecture 3.8 can be considered as a “generalization” of 2.5(ii).

It is easy to show that the cone  $C$  from 3.1 is a reflexive Gorenstein cone of index  $r$  if and only if  $\Delta = \Delta_1 + \dots + \Delta_r$  is a reflexive polytope. In this situation, we have

$$I_{\tilde{\Delta}} = t_{d+1} \cdots t_{d+r} S_{\tilde{\Delta}}.$$

Therefore one has canonical isomorphisms:

$$I_{\tilde{\Delta}}^k \cong S_{\tilde{\Delta}}^{\bar{k}}, \quad \forall k \in \mathbb{Z}_{>0}^r,$$

where the monomial basis in  $S_{\Delta}^{\bar{k}}$  can be identified with the set of all lattice points in  $\bar{k}_1\Delta_1 + \cdots + \bar{k}_r\Delta_r$ . The  $\bar{k}$ -homogeneous component of the corresponding toric residue map

$$\text{Res}_F^{\bar{k}} : S_{\Delta}^{\bar{k}} \rightarrow \mathbb{C}.$$

will be also called the  $\bar{k}$ -mixed toric residue.

#### 4. TORIC RESIDUE MIRROR CONJECTURE

Let  $M$  and  $N = \text{Hom}(M, \mathbb{Z})$  be the dual to each other abelian groups of rank  $d$ ,  $M_{\mathbb{R}}$  and  $N_{\mathbb{R}}$  their  $\mathbb{R}$ -scalar extensions and  $\Delta \subset M_{\mathbb{R}}$  a reflexive polytope with the unique interior lattice point  $0 \in M$ . Denote by  $\mathbb{P}_{\Delta}$  the Gorenstein toric Fano variety associated with  $\Delta$ . Let  $D_1, \dots, D_s$  be the toric divisors on  $\mathbb{P}_{\Delta}$  corresponding to the codimension-1 faces  $\Theta_1, \dots, \Theta_s$  and  $e_1, \dots, e_s$  the vertices of the dual reflexive polytope  $\Delta^* \subset N_{\mathbb{R}}$  such that

$$\Delta = \{x \in M_{\mathbb{R}} : \langle x, e_j \rangle \geq -1, \quad j = 1, \dots, s\},$$

$$\Theta_j = \Delta \cap \{x \in M_{\mathbb{R}} : \langle x, e_j \rangle = -1\}, \quad j \in \{1, \dots, s\}.$$

**Definition 4.1.** A Minkowski sum  $\Delta = \Delta_1 + \cdots + \Delta_r$  is called a *nef-partition* of the reflexive polytope  $\Delta$  if all vertices of  $\Delta_1, \dots, \Delta_r$  belong to  $M$ , and

$$\min_{x \in \Delta_i} \langle x, e_j \rangle \in \{0, -1\}, \quad \forall 1 \leq i \leq r, \quad \forall 1 \leq j \leq s.$$

Since  $\min_{x \in \Delta} \langle x, e_j \rangle = -1$  for all  $j \in \{1, \dots, s\}$ , the equality  $\min_{x \in \Delta_i} \langle x, e_j \rangle = -1$  holds exactly for one index  $i \in \{1, \dots, r\}$  if we fix a vertex  $e_j \in \Delta^*$ . Therefore, we can split the set of vertices  $\{e_1, \dots, e_s\} \subset \Delta^*$  into a disjoint union of subsets  $B_1, \dots, B_r$  where

$$B_i := \{e_j \in \{1, \dots, s\} : \min_{x \in \Delta_i} \langle x, e_j \rangle = -1\}.$$

Now we can define  $r$  nef Cartier divisors

$$E_i := \sum_{j: e_j \in B_i} D_j, \quad i = 1, \dots, r.$$

Therefore, a nef-partition  $\Delta = \Delta_1 + \cdots + \Delta_r$  of polytopes induces a partition of the anti-canonical divisor  $-K_{\mathbb{P}_{\Delta}} = D_1 + \cdots + D_n$  of  $\mathbb{P}_{\Delta}$  into a sum of  $r$  nef Cartier divisors:

$$-K_{\mathbb{P}_{\Delta}} = E_1 + \cdots + E_r.$$

Now it is easy to see that the above definition of the nef-partition is equivalent to the definition given in [Bo].

**Definition 4.2.** If  $\Delta = \Delta_1 + \cdots + \Delta_r$  is a nef-partition, then for any  $i = 1, \dots, r$  we denote

$$\nabla_i := \{y \in N_{\mathbb{R}} : \langle x, y \rangle \geq -\delta_{ij}, \ x \in \Delta_j, \ j = 1, \dots, r\}.$$

The lattice polytopes  $\nabla_1, \dots, \nabla_r$  define another nef-partition  $\nabla := \nabla_1 + \cdots + \nabla_r$  of the reflexive polytope  $\nabla \subset N_{\mathbb{R}}$  which is called the *dual nef-partition*.

The lattice polytopes  $\nabla_1, \dots, \nabla_r$  can be also defined as

$$\nabla_j := \text{conv}(\{0\} \cup B_j) \subset M_{\mathbb{R}}, \quad j = 1, \dots, r.$$

Moreover, one has two dual reflexive polytopes

$$\Delta^* = \text{conv}(\nabla_1 \cup \cdots \cup \nabla_r) \subset N_{\mathbb{R}}$$

$$\nabla^* = \text{conv}(\Delta_1 \cup \cdots \cup \Delta_r) \subset M_{\mathbb{R}}.$$

A nef-partition  $\Delta = \Delta_1 + \cdots + \Delta_r$  defines a family of  $(d-r)$ -dimensional Calabi-Yau complete intersections defined by vanishing of  $r$  Laurent polynomials  $f_1, \dots, f_r$  with Newton polytopes  $\Delta_1, \dots, \Delta_r$ . According to [Bo], the dual nef-partition  $\nabla = \nabla_1 + \cdots + \nabla_r$  defines the mirror dual family of Calabi-Yau complete intersections.

Define  $\mathcal{A}_j$  to be a subset in  $\Delta_j \cap M$  containing all vertices of  $\Delta_j$  and set  $A_j := \mathcal{A}_j \setminus \{0\}$ ,  $j = 1, \dots, r$ . It is easy to see that  $A_i \cap A_j = \emptyset$  for all  $i \neq j$ . We set

$$A_1 \cup \cdots \cup A_r := \{v_1, \dots, v_n\},$$

and define  $a_1, \dots, a_n \in \mathbb{C}$  to be the coefficients of the Laurent polynomials  $f_1, \dots, f_r$

$$f_j(t) := 1 - \sum_{i: v_i \in A_j} a_i t^{v_j}, \quad j = 1, \dots, r.$$

Let  $A := \{0\} \cup A_1 \cup \cdots \cup A_r$  and  $\tilde{\Delta} = \Delta_1 * \cdots * \Delta_r$  be the Cayley polytope. Denote by  $\pi$  the injective mapping

$$A_1 \cup \cdots \cup A_r \rightarrow \tilde{\Delta} \cap \tilde{M}$$

which sends a nonzero lattice point  $m \in A_j$  to  $(m, b_j)$  ( $1 \leq j \leq r$ ) and define

$$\tilde{A} := \pi(A_1 \cup \cdots \cup A_r) \cup \{(0, b_1), \dots, (0, b_r)\}.$$

We hold notations from [BM1, §4].

**Definition 4.3.** Choose a coherent triangulation  $\mathcal{T} = \{\tau_1, \dots, \tau_p\}$  of the reflexive polytope  $\nabla^* = \text{conv}(\Delta_1 \cup \cdots \cup \Delta_r)$  associated with  $A$  such that 0 is a vertex of all its  $d$ -dimensional simplices  $\tau_1, \dots, \tau_p$  and all elements of  $A$  appear as vertices of simplices in  $\mathcal{T}$ . Define a coherent triangulation  $\tilde{\mathcal{T}} = \{\tilde{\tau}_1, \dots, \tilde{\tau}_p\}$  of  $\tilde{\Delta} = \Delta_1 * \cdots * \Delta_r$  associated with  $\tilde{A}$  as follows: a  $(d+r-1)$ -dimensional simplex  $\tilde{\tau}_i \in \tilde{\mathcal{T}}$  is the convex hull of  $\pi$ -images of all nonzero vertices of  $\tau_i$  and  $\{(0, b_1), \dots, (0, b_r)\}$ . We call  $\tilde{\mathcal{T}}$  the **induced triangulation** of  $\tilde{\Delta}$ .

Let  $\mathbb{P} := \mathbb{P}_{\Sigma(\mathcal{T})}$  be the  $d$ -dimensional simplicial toric variety defined by the fan  $\Sigma(\mathcal{T}) \subset M_{\mathbb{R}}$  ( $\mathbb{P}$  is a partial crepant desingularization of the Gorenstein toric Fano variety  $\mathbb{P}_{\nabla}$ ) and denote by  $\mathbb{P}_{\beta}$  the *Morrison-Plesser moduli space* [BM1, Definition 3.3] corresponding to a lattice point

$$\beta = (\beta_1, \dots, \beta_n) \in R(\Sigma) = \{(x_1, \dots, x_n) \in \mathbb{Z}^n : x_1 v_1 + \dots + x_n v_n = 0\}$$

in the Mori cone  $K_{\text{eff}}(\mathbb{P})$ . One has a canonical surjective homomorphism

$$\psi_{\beta} : H^2(\mathbb{P}, \mathbb{Q}) \rightarrow H^2(\mathbb{P}_{\beta}, \mathbb{Q}).$$

**Definition 4.4.** By abuse of notations, let us denote by  $[D_j] \in H^2(\mathbb{P}_{\beta}, \mathbb{Q})$  ( $1 \leq j \leq n$ ) the image of  $[D_j] \in H^2(\mathbb{P}, \mathbb{Q})$  under  $\psi_{\beta}$ . Using the multiplication in the cohomology ring  $H^*(\mathbb{P}_{\beta}, \mathbb{Q})$ , we define the intersection product

$$\Phi_{\beta} := [E_1]^{(E_1, \beta)} \dots [E_r]^{(E_r, \beta)} \prod_{i: (D_i, \beta) < 0} [D_i]^{-(D_i, \beta) - 1}$$

considered as a cohomology class in  $H^{2(\dim \mathbb{P}_{\beta} - d)}(\mathbb{P}_{\beta}, \mathbb{Q})$  and call  $\Phi_{\beta}$  the **Morrison-Plesser class** corresponding to the nef-partition  $\Delta = \Delta_1 + \dots + \Delta_r$ .

**Definition 4.5.** Let  $k = (k_1, \dots, k_r) \in \mathbb{Z}_{>0}^r$  be a positive integral solution of

$$|k| = k_1 + \dots + k_r = d + r.$$

A polynomial  $P(x_1, \dots, x_n) \in \mathbb{Q}[x_1, \dots, x_n]$  is called  **$\bar{k}$ -homogeneous** if it is homogeneous of degree  $\bar{k}_i = k_i - 1$  with respect to every group of  $|A_i|$  variables  $x_j$  ( $v_j \in A_i$ ) ( $1 \leq i \leq r$ ).

Now we are able to formulate a generalized Toric Residue Mirror Conjecture:

**Conjecture 4.6.** Let  $\Delta = \Delta_1 + \dots + \Delta_r$  and  $\nabla = \nabla_1 + \dots + \nabla_r$  be two arbitrary dual nef-partitions. Choose any coherent triangulation  $\mathcal{T} = \{\tau_1, \dots, \tau_p\}$  of  $\nabla^*$  associated with  $A$  such that 0 is a vertex of all the simplices  $\tau_1, \dots, \tau_p$  as above. Then for any  $\bar{k}$ -homogeneous polynomial  $P(x_1, \dots, x_n) \in \mathbb{Q}[x_1, \dots, x_n]$  of degree  $d$  the Laurent expansion of the  $\bar{k}$ -mixed toric residue

$$R_P(a) := (-1)^d \text{Res}_F^{\bar{k}}(t_{d+1}^{\bar{k}_1} \dots t_{d+r}^{\bar{k}_r} P(a_1 t^{v_1}, \dots, a_n t^{v_n}))$$

at the vertex  $v_{\tilde{\mathcal{T}}} \in \text{Sec}(\tilde{\Delta})$  corresponding to the induced triangulation  $\tilde{\mathcal{T}} = \{\tilde{\tau}_1, \dots, \tilde{\tau}_p\}$  coincides with the generating function of intersection numbers

$$I_P(a) := \sum_{\beta \in K_{\text{eff}}(\mathbb{P})} I(P, \beta) a^{\beta},$$

where the sum runs over all integral points  $\beta = (\beta_1, \dots, \beta_n)$  of the Mori cone  $K_{\text{eff}}(\mathbb{P})$ ,  $a^{\beta} := a_1^{\beta_1} \dots a_n^{\beta_n}$ ,

$$I(P, \beta) = \int_{\mathbb{P}_{\beta}} P([D_1], \dots, [D_n]) \Phi_{\beta} = \langle P([D_1], \dots, [D_n]) \Phi_{\beta} \rangle_{\beta},$$

and  $\Phi_\beta \in H^{2(\dim \mathbb{P}_\beta - d)}(\mathbb{P}_\beta, \mathbb{Q})$  is the Morrison-Plesser class of  $\mathbb{P}_\beta$ . We assume  $I(P, \beta)$  to be zero if  $\mathbb{P}_\beta$  is empty.

## 5. COMPLETE INTERSECTIONS IN WEIGHTED PROJECTIVE SPACES

Let  $\mathbb{P} = \mathbb{P}(w_1, \dots, w_n)$  be a  $d$ -dimensional weighted projective space,  $n = d + 1$ . The fan  $\Sigma$  of  $\mathbb{P}(w_1, \dots, w_n)$  is determined by  $n$  vectors  $v_1, \dots, v_n \in M \simeq \mathbb{Z}^d$  which generate  $M$  and satisfy the relation

$$w_1 v_1 + \dots + w_n v_n = 0.$$

If we assume that  $\gcd(w_1, \dots, w_n) = 1$  and

$$w_i \mid (w_1 + \dots + w_n), \quad i = 1, \dots, n,$$

then  $\mathbb{P}$  is a Gorenstein toric Fano variety with the anticanonical divisor  $-K_{\mathbb{P}} = D_1 + \dots + D_n$ , where  $D_i$  is the toric divisor corresponding to the vector  $v_i$ . These divisors are related modulo rational equivalence as

$$\frac{[D_1]}{w_1} = \dots = \frac{[D_n]}{w_n} =: [D_0].$$

Consider a decomposition  $\{v_1, \dots, v_n\}$  into a disjoint union of  $r$  nonempty subsets  $A_1, \dots, A_r$  and define the divisors  $E_i := \sum_{j: v_j \in A_i} D_j$  on  $\mathbb{P}$  such that  $[E_i] = d_i [D_0]$ , where  $d_j = \sum_{i \in A_j} w_i$ ,  $j = 1, \dots, r$ . Note that the integers  $d_i$  satisfy

$d_1 + \dots + d_r = w_1 + \dots + w_n$ . Let  $\Delta_i := \text{conv}(\{0\} \cup A_i)$  ( $1 \leq i \leq r$ ). The polytopes  $\Delta_1, \dots, \Delta_r$  define a nef-partition  $\Delta := \Delta_1 + \dots + \Delta_r$  if and only if

$$w_i \mid d_j, \quad i = 1, \dots, n, \quad j = 1, \dots, r.$$

The following result generalizes [BM1, Theorem 7.3]:

**Theorem 5.1.** *Let  $P \in \mathbb{Q}[x_1, \dots, x_n]$  be a homogeneous polynomial of degree  $d$ . Then the generating function of intersection numbers on the Morrison-Plesser moduli spaces has the form*

$$I_P(y) = \nu \cdot P(w_1, \dots, w_n) \sum_{b \geq 0} \mu^b y^b = \frac{\nu \cdot P(w_1, \dots, w_n)}{1 - \mu y},$$

where

$$\nu := \frac{1}{w_1 \dots w_n}, \quad \mu := \frac{d_1^{d_1} \dots d_r^{d_r}}{w_1^{w_1} \dots w_n^{w_n}}, \quad y := a_1^{w_1} \dots a_n^{w_n}.$$

*Proof.* The lattice points  $\beta$  in the Mori cone of  $\mathbb{P}$  correspond to the linear relations  $bw_1 v_1 + \dots + bw_n v_n = 0$ ,  $b \in \mathbb{Z}_{\geq 0}$ . Therefore we set  $y := a_1^{w_1} \dots a_n^{w_n}$ .

The Morrison-Plesser moduli space  $\mathbb{P}_\beta$  is the  $(\sum_{i=1}^n w_i)b+d$ -dimensional weighted projective space:

$$\mathbb{P}(\underbrace{w_1, \dots, w_1}_{b+1}, \dots, \underbrace{w_n, \dots, w_n}_{b+1}).$$

It is easy to see that the Morrison-Plesser class defined by the nef-partition is

$$\Phi_\beta = (d_1[D_0])^{d_1 b} \dots (d_r[D_0])^{d_r b}.$$

Using  $\langle [D_0]^{\dim \mathbb{P}_\beta} \rangle_\beta = 1/w_1^{w_1 b+1} \dots w_n^{w_n b+1}$ , we obtain

$$\begin{aligned} I_P(y) &= \sum_{b \geq 0} \langle P([D_1], \dots, [D_n]) (d_1[D_0])^{d_1 b} \dots (d_r[D_0])^{d_r b} \rangle_\beta y^b \\ &= P(w_1, \dots, w_n) \sum_{b \geq 0} (d_1^{d_1} \dots d_r^{d_r})^b \langle [D_0]^{\dim \mathbb{P}_\beta} \rangle_\beta y^b \\ &= P(w_1, \dots, w_n) \sum_{b \geq 0} (d_1^{d_1} \dots d_r^{d_r})^b \frac{1}{w_1^{w_1 b+1} \dots w_n^{w_n b+1}} y^b \\ &= \nu \cdot P(w_1, \dots, w_n) \sum_{b \geq 0} \mu^b y^b \\ &= \frac{\nu \cdot P(w_1, \dots, w_n)}{1 - \mu y}. \end{aligned}$$

□

The convex hull of the vectors  $v_1, \dots, v_n$  is a reflexive polytope (simplex)  $\nabla^* \subset M_\mathbb{R} \cong \mathbb{R}^d$ . Let  $\widetilde{M} := M \oplus \mathbb{Z}^r$  be an extension of the lattice  $M$  and  $\{b_1, \dots, b_r\}$  the standard basis of  $\mathbb{Z}^r$ . The  $(d+r-1)$ -dimensional Cayley polytope

$$\widetilde{\Delta} = \Delta_1 * \dots * \Delta_r$$

is the convex hull of  $(d+r+1)$  points:  $e_1 = (0, b_1), \dots, e_r = (0, b_r)$  and  $u_k = (v_k, b_j)$  ( $k = 1, \dots, d+1$ ), where  $v_k \in A_j$ . We denote this set of points by  $\widetilde{A}$ . The points from  $\widetilde{A}$  are affinely dependent, while any proper subset of  $\widetilde{A}$  is affinely independent, i.e., defines a circuit (see [GKZ, Chapter 7]). It is easy to see that the only affine relation (up to a real multiple) between the points from  $\widetilde{A}$  is

$$d_1 e_1 + \dots + d_r e_r - w_1 u_1 - \dots - w_n u_n = 0.$$

Thus by [GKZ, Chapter 7, Proposition 1.2], polytope  $\widetilde{\Delta}$  has exactly two triangulations: the triangulation  $\mathcal{T} = \mathcal{T}_1$  with the simplices  $\text{conv}(\widetilde{A} \setminus \{e_i\})$ ,  $i = 1, \dots, r$ , and the triangulation  $\mathcal{T}_2$  with the simplices  $\text{conv}(\widetilde{A} \setminus \{u_k\})$ ,  $k = 1, \dots, n$ . Note that

$$(1) \quad \text{Vol}(\text{conv}(\widetilde{A} \setminus \{e_i\})) = d_i, \quad i = 1, \dots, r,$$

$$(2) \quad \text{Vol}(\text{conv}(\widetilde{A} \setminus \{u_k\})) = w_k, \quad k = 1, \dots, n.$$

Therefore  $\text{Vol}(\tilde{\Delta}) = \sum_{i=1}^r d_i = \sum_{k=1}^n w_k$ .

Let

$$f_j(t) := 1 - \sum_{i: v_i \in A_j} a_i t^{v_i} \in \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}], \quad j = 1, \dots, r$$

be generic Laurent polynomials. Denote by

$$F(t) = t_{d+1} f_1(t) + \dots + t_{d+r} f_r(t)$$

a Laurent polynomial whose support polytope is  $\tilde{\Delta}$ .

The next statement follows directly from [GKZ, Chapter 9, Proposition 1.8] and from the equalities (1), (2).

**Proposition 5.2.** *The  $\tilde{A}$ -discriminant of  $F$  is equal (up to sign) to the binomial*

$$D_{\tilde{A}}(F) = \prod_{k=1}^n w_k^{w_k} - \prod_{i=1}^r d_i^{d_i} \prod_{k=1}^n a_k^{w_k} = \prod_{k=1}^n w_k^{w_k} (1 - \mu y),$$

where  $y = \prod_{k=1}^n a_k^{w_k}$  and the first summand in  $D_{\tilde{A}}(F)$  corresponds to the triangulation  $\mathcal{T}$ .

**Theorem 5.3.** *Let  $P(x_1, \dots, x_n) \in \mathbb{C}[x_1, \dots, x_n]$  be a  $\bar{k}$ -homogeneous polynomial with  $|\bar{k}| = d$ . Then*

$$R_P(a) = (-1)^d \text{Res}_F^{\bar{k}} \left( t_{d+1}^{\bar{k}_1} \dots t_{d+r}^{\bar{k}_r} P(a_1 t^{v_1}, \dots, a_n t^{v_n}) \right) = \frac{\nu \cdot P(w_1, \dots, w_n)}{1 - \mu y},$$

where  $y := a_1^{w_1} \dots a_n^{w_n}$ .

*Proof.* By Proposition 2.6 the toric residue  $R_P(a)$  is the following sum over the critical points  $\xi$  of the polynomial  $F_1(t, y) := f_1(t) + y_2 f_2(t) + \dots + y_r f_r(t)$ , where  $(t, y) \in (\mathbb{C}^*)^d \times (\mathbb{C}^*)^{r-1}$ :

$$R_P(a) = (-1)^d \sum_{\xi \in V_F} \frac{P(a_1 \xi^{v_1}, \dots, a_n \xi^{v_n})}{F_1(\xi) H_{F_1}^1(\xi)}.$$

We rewrite the polynomial  $F_1$  as

$$F_1 = y_2 + \dots + y_r + 1 - \sum_{i=1}^n c_i t^{v_i},$$

where  $c_i = y_j \cdot a_i$  if  $v_i \in A_j$ . Then at the critical point  $\xi$ , we have

$$c_1 \frac{\xi^{v_1}}{w_1} = \dots = c_n \frac{\xi^{v_n}}{w_n} = z$$

and

$$z^{w_1 + \dots + w_n} = \left( \frac{c_1}{w_1} \right)^{w_1} \dots \left( \frac{c_n}{w_n} \right)^{w_n}.$$



Moreover, at the critical points one has:

$$f_2(\xi) = \cdots = f_r(\xi) = 0,$$

which is equivalent to

$$f_j(\xi) = 1 - \sum_{i: v_i \in A_j} a_i \xi^{v_i} = 1 - \left( \sum_{i: v_i \in A_j} w_i \right) \frac{z}{\eta_j} = 1 - \frac{d_j z}{\eta_j} = 0, \quad j = 2, \dots, r,$$

where  $\eta_j$  is the value of  $y_j$  at the critical point. Hence, it is easy to see that  $\eta_j = d_j z$ , ( $j = 2, \dots, r$ ) and  $F = 1 - d_1 z$ , which implies

$$(3) \quad z^{w_1 + \cdots + w_n} = \left( \frac{a_1}{w_1} \right)^{w_1} \cdots \left( \frac{a_n}{w_n} \right)^{w_n} d_2^{d_2} \cdots d_r^{d_r} \cdot z^{d_2 + \cdots + d_r},$$

or, equivalently,

$$z^{d_1} = \left( \frac{a_1}{w_1} \right)^{w_1} \cdots \left( \frac{a_n}{w_n} \right)^{w_n} d_2^{d_2} \cdots d_r^{d_r}.$$

The value of the Hessian  $H_F^1$  at  $\xi$  equals

$$H_F^1(\xi) = (-1)^d w_1 \cdots w_n d_1 \cdots d_r z^{d+r-1}.$$

Since there are exactly  $w_1 + \cdots + w_n = d_1 + \cdots + d_r$  critical points of  $F$ , the summation over the critical points is equivalent to the summation over the roots of (3), we get

$$\begin{aligned} R_P(y) &= \sum_{z^{d_1} = \left( \frac{a_1}{w_1} \right)^{w_1} \cdots \left( \frac{a_n}{w_n} \right)^{w_n} d_2^{d_2} \cdots d_r^{d_r}} \frac{P(w_1, \dots, w_n)}{w_1 \cdots w_n d_1 (1 - d_1 z)} \\ &= \frac{P(w_1, \dots, w_n)}{w_1 \cdots w_n} \sum_{b \geq 0} (d_1^{d_1} \cdots d_r^{d_r})^b \left( \frac{a_1^{w_1} \cdots a_n^{w_n}}{w_1^{w_1} \cdots w_n^{w_n}} \right)^b \\ &= \frac{\nu \cdot P(w_1, \dots, w_n)}{1 - \mu y}. \end{aligned}$$

□

## 6. COMPLETE INTERSECTIONS IN PRODUCT OF PROJECTIVE SPACES

In this section we check the Toric Residue Mirror Conjecture for nef-partitions corresponding to mirrors of complete intersections in a product of projective spaces  $\mathbb{P} = \mathbb{P}^{d_1} \times \cdots \times \mathbb{P}^{d_p}$  of dimension  $d = d_1 + \cdots + d_p$ . We set  $n_i := d_i + 1$  and denote by  $N = (n_{ij})$  an integral  $p \times r$ -matrix with non-negative elements having columns  $\mathbf{n}_1, \dots, \mathbf{n}_r \in \mathbb{Z}_{\geq 0}^p$ . A complete intersection  $V$  of  $r$  hypersurfaces  $V_1, \dots, V_r$  in  $\mathbb{P}$  of

multidegrees  $\mathbf{n}_1, \dots, \mathbf{n}_r$  is a Calabi-Yau  $(d-r)$ -fold if and only if  $\sum_{j=1}^r n_{ij} = n_i$  ( $i = 1, \dots, p$ ). We will use the standard notation

$$\left( \begin{array}{c|ccc} \mathbb{P}^{d_1} & n_{11} & \cdots & n_{1r} \\ \vdots & \vdots & & \vdots \\ \mathbb{P}^{d_p} & n_{p1} & \cdots & n_{pr} \end{array} \right)$$

to denote this complete intersection.

The cone of effective curves  $K_{\text{eff}}(\mathbb{P})$  is isomorphic to  $\mathbb{R}_{\geq 0}^p$  and its integral part  $K_{\text{eff}}(\mathbb{P})_{\mathbb{Z}}$  consists of the points  $\beta = (b_1, \dots, b_p) \in \mathbb{Z}_{\geq 0}^p$ . Thus, the Morrison-Plesser moduli spaces are the products of projective spaces:  $\mathbb{P}_{\beta} = \mathbb{P}^{n_1 b_1 + d_1} \times \cdots \times \mathbb{P}^{n_p b_p + d_p}$  and the generating function for intersection numbers may be written

$$I_P(y) = \sum_{b_1, \dots, b_p \geq 0} I(P, \beta) y_1^{b_1} \cdots y_p^{b_p}.$$

**Theorem 6.1.** *The generating function for intersection numbers associated with the monomial  $x^k = x_1^{k_1} \cdots x_p^{k_p}$  can be written as the integral*

$$(4) \quad I_{x^k}(y) = \left( \frac{1}{2\pi i} \right)^p \int_{\Gamma} \frac{z_1^{k_1} \cdots z_p^{k_p} dz_1 \wedge \cdots \wedge dz_p}{G_1(z) \cdots G_p(z)},$$

where the polynomials  $G_i$  have the form

$$G_i = z_i^{n_i} - \prod_{j=1}^r (n_{1j} z_1 + \cdots + n_{pj} z_p)^{n_{ij}} y_i, \quad i = 1, \dots, p,$$

and  $\Gamma$  is the compact cycle in  $\mathbb{C}^p$  defined by  $\Gamma = \{|G_1| = \cdots = |G_p| = \delta\}$  for small positive  $\delta$ .

*Proof.* Let  $[H_i]$  denote the class of a hyperplane section in  $\mathbb{P}^{d_i}$ . The class of the divisor  $E_j$  defining hypersurface  $V_j$  equals

$$[E_j] = n_{1j}[H_1] + \cdots + n_{pj}[H_p], \quad j = 1, \dots, r.$$

Hence, the coefficients of the series  $I_{x^k}(y)$  are

$$\langle [H_1]^{k_1} \cdots [H_p]^{k_p} \prod_{j=1}^r (n_{1j}[H_1] + \cdots + n_{pj}[H_p])^{n_{1j}b_1 + \cdots + n_{pj}b_p} \rangle_{\beta}.$$

The lattice points  $\beta = (b_1, \dots, b_p) \in \mathbb{Z}_{\geq 0}^p$  in the integral part of the Mori cone  $K_{\text{eff}}(\mathbb{P})$  correspond to the  $p$  linear relations

$$b_i v_{i1} + \cdots + b_i v_{in_i} = 0, \quad i = 1, \dots, p,$$

where  $v_{j1}, \dots, v_{jn_j}$  generate a lattice  $M_j$  of rank  $d_j$  ( $1 \leq j \leq p$ ). Therefore we set  $y_i := a_{i1} \cdots a_{in_i}$ . Using the property of the integral

$$\left(\frac{1}{2\pi i}\right)^p \int_{\gamma_\rho} z_1^{m_1-1} \cdots z_p^{m_p-1} dz = \begin{cases} 1, & m_1 = \cdots = m_p = 0, \\ 0, & \text{otherwise,} \end{cases}$$

where  $\gamma_\rho = \{|z_1| = \cdots = |z_p| = \rho\}$  is the cycle winding around the origin ( $\rho > 0$  is small) and the fact that the intersection numbers on  $\mathbb{P}$  are

$$\langle [H_1]^{l_1} \cdots [H_p]^{l_p} \rangle_\beta = \begin{cases} 1, & l_j = n_j b_j + d_j, \quad j = 1, \dots, r, \\ 0, & \text{otherwise,} \end{cases}$$

we can represent the functions  $I(x^k, \beta)$  by integrals

$$I(x^k, \beta) = \left(\frac{1}{2\pi i}\right)^p \int_{\gamma_\rho} \frac{z_1^{k_1} \cdots z_p^{k_p} \prod_{j=1}^r (n_{1j} z_1 + \cdots + n_{pj} z_p)^{n_{1j} b_1 + \cdots + n_{pj} b_p} dz}{z_1^{n_1 b_1 + d_1 + 1} \cdots z_p^{n_p b_p + d_p + 1}}.$$

Denote

$$F_i(z) := \prod_{j=1}^r (n_{1j} z_1 + \cdots + n_{pj} z_p)^{n_{ij}}, \quad i = 1, \dots, p.$$

If  $z \in \gamma_\rho$  for some fixed  $\rho$ , then the geometric series

$$z_1^{k_1 - n_1} \cdots z_p^{k_p - n_p} \sum_{b_1, \dots, b_p \geq 0} \left(\frac{F_1(z)y_1}{z_1^{n_1}}\right)^{b_1} \cdots \left(\frac{F_p(z)y_p}{z_p^{n_p}}\right)^{b_p} = \frac{z_1^{k_1} \cdots z_p^{k_p}}{\prod_{i=1}^p [z_i^{n_i} - F_i(z)y_i]}$$

converges absolutely and uniformly for all  $y$  from the neighborhood  $\mathcal{U}_\varepsilon = \{y : \|y\| < \varepsilon\}$ , where  $0 < \varepsilon < \min_{i=1, \dots, p} (\rho^{n_i} / M_i)$ ,  $M_i = \max_{z \in \gamma_\rho} |F_i(z)|$ . Integrating the last expression and changing the order of integration and summation, we get

$$I_{x^k}(y) = \left(\frac{1}{2\pi i}\right)^p \int_{\gamma_\rho} \frac{z_1^{k_1} \cdots z_p^{k_p} dz_1 \wedge \cdots \wedge dz_p}{\prod_{i=1}^p [z_i^{n_i} - F_i(z)y_i]}.$$

For fixed  $y \in \mathcal{U}_\varepsilon$ , using the Rouché's principle for residues (see [Ts, Chapter 2, §8] or [AY, Lemma 4.9]), the cycle  $\gamma_\rho$  can be replaced by another homologically equivalent cycle  $\Gamma$  of the following form:

$$\gamma_\rho \sim \Gamma = \{z : |z_1^{n_1} - F_1(z)y_1| = \cdots = |z_p^{n_p} - F_p(z)y_p| = \delta\}$$

for some  $\delta > 0$ . Therefore, we have

$$I_{x^k}(y) = \left(\frac{1}{2\pi i}\right)^p \int_{\Gamma} \frac{z_1^{k_1} \cdots z_p^{k_p} dz_1 \wedge \cdots \wedge dz_p}{\prod_{i=1}^p [z_i^{n_i} - F_i(z)y_i]}$$

which finishes the proof.  $\square$

The Conjecture 4.6 follows now from a general result in [BM2] which identifies

$$\left(\frac{1}{2\pi i}\right)^p \int_{\Gamma} \frac{z_1^{k_1} \cdots z_p^{k_p} dz_1 \wedge \cdots \wedge dz_p}{G_1(z) \cdots G_p(z)}$$

with the toric residue.

## 7. COMPUTATION OF YUKAWA $(d-r)$ -POINT FUNCTIONS

Let  $\Delta = \Delta_1 + \cdots + \Delta_r$  be a nef-partition of a reflexive polytope  $\Delta$ ,  $A_i \subset \partial \nabla^* \cap \Delta_i \cap M$  a subset containing all nonzero vertices of  $\Delta_i$  ( $1 \leq i \leq r$ ). We set  $A_1 \cup \cdots \cup A_r := \{v_1, \dots, v_n\}$  and consider a  $\Delta_1 * \cdots * \Delta_r$ -regular sequence of Laurent polynomials

$$f_j(t) := 1 - \sum_{i: v_i \in A_j} a_i t^{v_i} \in \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}], \quad j = 1, \dots, r,$$

which define  $r$  affine hypersurfaces

$$Z_{f_j} := \{t \in \mathbb{T} \cong (\mathbb{C}^*)^d : f_j(t) = 0\}, \quad j = 1, \dots, r,$$

The compactification  $\overline{Z}_f$  in  $\mathbb{P}_{\Delta}$  of the affine complete intersection  $Z_f := Z_{f_1} \cap \cdots \cap Z_{f_r}$  is a  $(d-r)$ -dimensional projective Calabi-Yau variety with at worst Gorenstein canonical singularities. Using the Poincaré residue mapping

$$\mathbf{Res} : H^d(\mathbb{T} \setminus Z_{f_1} \cup \cdots \cup Z_{f_r}) \rightarrow H^{d-r}(Z_{f_1} \cap \cdots \cap Z_{f_r})$$

one can construct a nowhere vanishing section of the canonical bundle of  $\overline{Z}_f$  as

$$\Omega := \mathbf{Res} \left( \frac{1}{f_1 \cdots f_r} \frac{dt_1}{t_1} \wedge \cdots \wedge \frac{dt_d}{t_d} \right).$$

**Definition 7.1.** Let  $Q(x_1, \dots, x_n) \in \mathbb{Q}[x_1, \dots, x_n]$  be a homogeneous polynomial of degree  $d-r$ . The  $Q$ -Yukawa  $(d-r)$ -point function is defined by the formula

$$Y_Q(a_1, \dots, a_n) := \frac{(-1)^{\frac{(d-r)(d-r-1)}{2}}}{(2\pi i)^{d-r}} \int_{Z_f} \Omega \wedge Q \left( a_1 \frac{\partial}{\partial a_1}, \dots, a_n \frac{\partial}{\partial a_n} \right) \Omega,$$

where the differential operators  $a_1 \partial / \partial a_1, \dots, a_n \partial / \partial a_n$  are determined by the Gauß-Manin connection. If  $\overline{k} = (\overline{k}_1, \dots, \overline{k}_r)$  is a nonnegative integral vector with  $|\overline{k}| = d-r$  and  $Q(x_1, \dots, x_n)$  is a  $\overline{k}$ -homogeneous polynomial ( $\deg x_j = k_i \Leftrightarrow v_j \in A_i$ ), then

$$Q \left( a_1 \frac{\partial}{\partial a_1}, \dots, a_n \frac{\partial}{\partial a_n} \right) \Omega = (-1)^{d-r} \mathbf{Res} \left( \frac{Q(a_1 t^{v_1}, \dots, a_n t^{v_n})}{f_1^{\overline{k}_1+1} \cdots f_r^{\overline{k}_r+1}} \frac{dt_1}{t_1} \wedge \cdots \wedge \frac{dt_d}{t_d} \right).$$

**Theorem 7.2.** *Let  $Q(x_1, \dots, x_n) \in \mathbb{C}[x_1, \dots, x_n]$  be a  $\bar{k}$ -homogeneous polynomial with  $|\bar{k}| = d - r$ . We define*

$$P(x_1, \dots, x_n) := \prod_{j=1}^r \left( \sum_{i: v_i \in A_j} x_i \right) Q(x_1, \dots, x_n).$$

*Then the Yukawa  $(d - r)$ -point function is equal to the  $k$ -mixed toric residue*

$$Y_Q(a_1, \dots, a_n) = (-1)^d \text{Res}_F^k \left( t_{d+1}^{\bar{k}_1+1} \cdots t_{d+r}^{\bar{k}_r+1} P(a_1 t^{v_1}, \dots, a_n t^{v_n}) \right).$$

*Proof.* We sketch only the idea of the proof. The hypersurface

$$Z_F = \{t_{d+1}f_1 + \cdots + t_{d+r}f_r = 1\}$$

in  $(\mathbb{C}^*)^d \times \mathbb{C}^r$  is a  $\mathbb{C}^{r-1}$ -bundle over  $(\mathbb{C}^*)^d \setminus (Z_{f_1} \cap \cdots \cap Z_{f_r})$ . This fact allows one to identify the primitive parts of the cohomology groups  $H^{d-r}(Z_{f_1} \cap \cdots \cap Z_{f_r})$  and  $H^{d-1}(Z_F)$  together with their intersection forms. By the result of Mavlyutov [Mav2], one can compute the intersection form on  $H^{d-1}(Z_F)$  using toric residues.  $\square$

**Example 7.3.** Consider the mirror family  $V^*$  to Calabi-Yau complete intersections  $V$  of  $r$  hypersurfaces of degrees  $d_1, \dots, d_r$  respectively in  $\mathbb{P}^d$ ,  $d_1 + \cdots + d_r = d + 1$ . Its nef-partition can be constructed as follows. Let  $v_1, \dots, v_d$  be a basis vectors of the lattice  $M$  and

$$v_{d+1} := -v_1 - \cdots - v_d.$$

We divide the set  $\{v_1, \dots, v_{d+1}\}$  into a disjoint union of  $r$  subsets  $A_1, \dots, A_r$  such that  $|A_i| = d_i$ . For  $j = 1, \dots, r$ , we define Laurent polynomials

$$f_j(t) := 1 - \sum_{i: v_i \in A_j} a_i t^{v_i} \in \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}].$$

Then the affine part of  $V^*$  is the complete intersection  $Z_f \subset \mathbb{T}$  of hypersurfaces  $Z_{f_1}, \dots, Z_{f_r} \subset \mathbb{T}$  defined by polynomials  $f_1, \dots, f_r$ . The Yukawa coupling for  $V^*$  has been computed in [BvS, Proposition 5.1.2]:

$$Y_Q(y) = \frac{d_1 \cdots d_r Q(1, \dots, 1)}{1 - \mu y},$$

where  $y = a_1 \cdots a_n$  and  $\mu = \prod_{i=1}^r d_i^{d_i}$ .

**Example 7.4.** Consider Calabi-Yau varieties  $V$  obtained as the complete intersection of hypersurfaces  $V_1, V_2, V_3$  in  $\mathbb{P}^3 \times \mathbb{P}^3$  of degrees  $(3, 0)$ ,  $(0, 3)$  and  $(1, 1)$  respectively of the type

$$\left( \begin{array}{c} \mathbb{P}^3 \\ \mathbb{P}^3 \end{array} \middle| \begin{array}{ccc} 3 & 0 & 1 \\ 0 & 3 & 1 \end{array} \right).$$

Let  $M \cong \mathbb{Z}^6$  and  $\nabla^* = \text{conv}(\Delta_1 \cup \Delta_2 \cup \Delta_3) \subset M_{\mathbb{R}}$  be a reflexive polytope defined by the polytopes  $\Delta_1 := \text{conv}\{0, v_1, v_2, v_3\}$ ,  $\Delta_2 := \text{conv}\{0, v_5, v_6, v_7\}$  and  $\Delta_3 := \text{conv}\{0, v_4, v_8\}$ , where

$$\begin{aligned} v_1 &= (1, 0, 0, 0, 0, 0), v_2 = (0, 1, 0, 0, 0, 0), v_3 = (0, 0, 1, 0, 0, 0), \\ v_4 &= (-1, -1, -1, 0, 0, 0), v_5 = (0, 0, 0, 1, 0, 0), v_6 = (0, 0, 0, 0, 1, 0), \\ v_7 &= (0, 0, 0, 0, 0, 1), v_8 = (0, 0, 0, -1, -1, -1). \end{aligned}$$

The nef-partition  $\Delta = \Delta_1 + \Delta_2 + \Delta_3$  corresponds to mirrors  $V^*$  of  $V = V_1 \cap V_2 \cap V_3$ . We define the disjoint sets:  $A_1 := \{v_1, v_2, v_3\}$ ,  $A_2 := \{v_5, v_6, v_7\}$ ,  $A_3 := \{v_4, v_8\}$  and the Laurent polynomials

$$\begin{aligned} f_1(t) &:= 1 - \sum_{i: v_i \in A_1} a_i t^{v_i} = 1 - a_1 t_1 - a_2 t_2 - a_3 t_3, \\ f_2(t) &:= 1 - \sum_{i: v_i \in A_2} a_i t^{v_i} = 1 - a_5 t_4 - a_6 t_5 - a_7 t_6, \\ f_3(t) &:= 1 - \sum_{i: v_i \in A_3} a_i t^{v_i} = 1 - a_4 t_1^{-1} t_2^{-1} t_3^{-1} - a_8 t_4^{-1} t_5^{-1} t_6^{-1}. \end{aligned}$$

The complete intersection  $Z_f := Z_{f_1} \cap Z_{f_2} \cap Z_{f_3}$  of the affine hypersurfaces

$$Z_{f_j} = \{t \in (\mathbb{C}^*)^6 : f_j(t) = 0\}, \quad j = 1, 2, 3$$

is an affine part of  $V^*$ .

Denote by  $y_1 = 3^3 a_1 a_2 a_3 a_4$ ,  $y_2 = 3^3 a_5 a_6 a_7 a_8$  the new variables and by  $\theta_1 := y_1 \partial / \partial y_1$ ,  $\theta_2 := y_2 \partial / \partial y_2$  the corresponding logarithmic partial derivations. Given a form-residue

$$\Omega := \mathbf{Res} \left( \frac{1}{f_1 f_2 f_3} \frac{dt_1}{t_1} \wedge \cdots \wedge \frac{dt_6}{t_6} \right) \in H^3(Z_f),$$

we compute the 2-parameter Yukawa couplings defined as integrals

$$Y^{(k_1, k_2)}(y_1, y_2) = \frac{-1}{(2\pi i)^3} \int_{Z_f} \Omega \wedge \theta_1^{k_1} \theta_2^{k_2} \Omega, \quad k_1 + k_2 = 3.$$

**Proposition 7.5.** *The Yukawa couplings are*

$$\begin{aligned} Y^{(3,0)}(y_1, y_2) &= \frac{9y_1}{(1 - y_1 - y_2)(1 - y_1)^2}, \quad Y^{(2,1)}(y_1, y_2) = \frac{9}{(1 - y_1 - y_2)(1 - y_1)}, \\ Y^{(1,2)}(y_1, y_2) &= \frac{9}{(1 - y_1 - y_2)(1 - y_2)}, \quad Y^{(0,3)}(y_1, y_2) = \frac{9y_2}{(1 - y_1 - y_2)(1 - y_2)^2}. \end{aligned}$$

*Remark 7.6.* Note that the functions in Proposition 7.5 are completely consistent with Yukawa couplings from [BvS, §8.3]. Indeed, if we put  $K(y_1, y_2) = Y^{(3,0)} +$

$3Y^{(2,1)} + 3Y^{(1,2)} + Y^{(0,3)}$  and consider restriction to the diagonal subfamily  $y = y_1 = y_2$ , then we get the same expression as in [BvS]:

$$K(y, y) = \frac{18(3-2y)}{(1-2y)(1-y)^2}.$$

Denote by  $F(t) := t_7 f_1(t) + t_8 f_2(t) + t_9 f_3(t)$  the Cayley polynomial associated with Laurent polynomials  $f_1, f_2, f_3$ , and by  $\tilde{\Delta} = \Delta_1 * \Delta_2 * \Delta_3 \subset \tilde{M}_{\mathbb{R}} = M_{\mathbb{R}} \oplus \mathbb{R}^3$  its supporting polytope which is the Cayley polytope associated with  $\Delta_1, \Delta_2, \Delta_3$ .

**Proposition 7.7.** *Let  $A := \tilde{\Delta} \cap \tilde{M}$  and  $F(t)$  be the Cayley polynomial as above. Then the principal  $A$ -determinant of  $F$  has the form*

$$E_A(F) = (a_1 \cdots a_8)^{12} (1-y_1)^3 (1-y_1)^3 (1-y_1-y_2).$$

*Remark 7.8.* It is easy to see that the products of  $Y^{(k_1, k_2)}$  by  $E_A(F)$  are polynomials in  $a_1, \dots, a_8$ .

Let us find the generating function  $I_P(y)$  for the monomial  $P(x) = x_1^{k_1} x_2^{k_2}$ . There are two linear independent integral relations between  $v_1, \dots, v_8$ :

$$v_1 + \cdots + v_4 = 0, \quad v_5 + \cdots + v_8 = 0.$$

Hence the Mori cone  $K_{\text{eff}}(\mathbb{P})$  is spanned by the vectors

$$l^{(1)} = (1, 1, 1, 1, 0, 0, 0, 0), \quad l^{(2)} = (0, 0, 0, 0, 1, 1, 1, 1)$$

and the Morrison-Plesser moduli spaces are  $\mathbb{P}_{\beta} = \mathbb{P}^{4b_1+3} \times \mathbb{P}^{4b_2+3}$  ( $b_1, b_2 \in \mathbb{Z}_{\geq 0}$ ). The cohomology ring of  $\mathbb{P}_{\beta}$  is generated by two hyperplane classes:  $[H_1]$  and  $[H_2]$ . We set  $E_1 := 3[H_1]$ ,  $E_2 := 3[H_2]$  and  $E_3 := [H_1] + [H_2]$ . Then the nef-partition of the anticanonical divisor  $-K_{\mathbb{P}}$  corresponding to the nef-partition  $\Delta = \Delta_1 + \Delta_2 + \Delta_3$  is defined by  $-K_{\mathbb{P}} = E_1 + E_2 + E_3$ . Therefore, the Morrison-Plesser cohomology class associated with the nef-partition of  $-K_{\mathbb{P}}$  equals

$$\Phi_{\beta} = (3[H_1])^{3b_1} (3[H_2])^{3b_2} ([H_1] + [H_2])^{b_1+b_2}$$

and the generating function for intersection numbers can be written

$$I_P(y) = \sum_{b_1, b_2 \geq 0} \langle [H_1]^{k_1+3b_1+1} [H_2]^{k_2+3b_2+1} ([H_1] + [H_2])^{b_1+b_2+1} \rangle_{\beta} y_1^{b_1} y_2^{b_2}.$$

Intersection theory on  $\mathbb{P}_{\beta}$  implies

$$I_P(y) = 9 \sum_{b_1, b_2 \geq 0} \frac{(b_1 + b_2 + 1)!}{(b_1 - k_1 + 2)!(b_2 - k_2 + 2)!} y_1^{b_1} y_2^{b_2}.$$

By Theorem 6.1 we can write  $I_P(y)$  as the integral

$$I_P(y) = \frac{1}{(2\pi i)^2} \int_{\Gamma} \frac{9z_1^{k_1+1} z_2^{k_2+1} (z_1 + z_2) dz_1 \wedge dz_2}{(z_1^4 - z_1^3(z_1 + z_2)y_1)(z_2^4 - z_2^3(z_1 + z_2)y_2)}$$

with the cycle  $\Gamma = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1^4 - z_1^3(z_1 + z_2)y_1| = |z_2^4 - z_2^3(z_1 + z_2)y_2| = \delta\}$ ,  $\delta > 0$ . Computing the last integrals, we get the same rational functions as in Proposition 7.5.

**Example 7.9.** Consider Calabi-Yau varieties  $V$  obtained as complete intersections of two hypersurfaces of degrees  $(4, 0)$ ,  $(1, 2)$  in  $\mathbb{P} = \mathbb{P}^4 \times \mathbb{P}^1$  which corresponds to the configuration

$$\left( \begin{array}{c|cc} \mathbb{P}^4 & 4 & 1 \\ \hline \mathbb{P}^1 & 0 & 2 \end{array} \right).$$

This example was investigated in details by Hosono, Klemm, Theisen and Yau (cf. [HKTY]). The corresponding nef-partition  $\Delta = \Delta_1 + \Delta_2 \subset M_{\mathbb{R}} \cong \mathbb{R}^5$  consists of polytopes  $\Delta_1 := \text{conv}\{0, v_1, v_2, v_3, v_4\}$  and  $\Delta_2 := \text{conv}\{0, v_5, v_6, v_7\}$ , where

$$\begin{aligned} v_1 &= (1, 0, 0, 0, 0), & v_2 &= (0, 1, 0, 0, 0), & v_3 &= (0, 0, 1, 0, 0), & v_4 &= (0, 0, 0, 1, 0), \\ v_5 &= (-1, -1, -1, -1, 0), & v_6 &= (0, 0, 0, 0, 1), & v_7 &= (0, 0, 0, 0, -1). \end{aligned}$$

We have two disjoint sets:  $A_1 := \{v_1, v_2, v_3, v_4\}$  and  $A_2 := \{v_5, v_6, v_7\}$  which are the vertices of the reflexive polytope  $\nabla^*$  and define the Laurent polynomials

$$\begin{aligned} f_1(t) &= 1 - \sum_{i: v_i \in A_1} a_i t^{v_i} = 1 - a_1 t_1 - a_2 t_2 - a_3 t_3 - a_4 t_4, \\ f_2(t) &= 1 - \sum_{i: v_i \in A_2} a_i t^{v_i} = 1 - a_5 (t_1 t_2 t_3 t_4)^{-1} - a_6 t_5 - a_7 t_5^{-1}. \end{aligned}$$

Denote by  $y_1 := a_1 \cdots a_5$ ,  $y_2 := a_6 a_7$  the new variables and  $\theta_1 := y_1 \partial / \partial y_1$ ,  $\theta_2 := y_2 \partial / \partial y_2$  the corresponding logarithmic partial derivations. Let  $\Omega$  be a form defined by

$$\Omega := \mathbf{Res} \left( \frac{1}{f_1 f_2} \frac{dt_1}{t_1} \wedge \frac{dt_2}{t_2} \right) \in H^3(Z_{f_1} \cap Z_{f_2}).$$

Then the Yukawa coupling associated with  $f_1, f_2$  is the integral

$$Y^{(k_1, k_2)}(y_1, y_2) := \frac{-1}{(2\pi i)^3} \int_{Z_f} \Omega \wedge \theta_1^{k_1} \theta_2^{k_2} \Omega, \quad k_1 + k_2 = 3,$$

where  $Z_f := Z_{f_1} \cap Z_{f_2}$  is an affine Calabi-Yau complete intersection which compactification forms a mirror dual family to  $V$ .

**Proposition 7.10.** [HKTY] *The Yukawa couplings  $Y^{(k_1, k_2)}(y)$  are:*

$$\begin{aligned} Y^{(3,0)}(y) &= \frac{8}{D_0}, & Y^{(2,1)}(y) &= \frac{4(1 - 256y_1 + 4y_2)}{D_0 D_1}, \\ Y^{(1,2)}(y) &= \frac{8y_2(3 - 512y_1 + 4y_2)}{D_0 D_1^2}, \\ Y^{(0,3)}(y) &= \frac{4y_2(1 - 256y_1 + 24y_2 - 3072y_1 y_2 + 16y_2^2)}{D_0 D_1^3}, \end{aligned}$$



where

$$D_0 := (1 - 256y_1)^2 - 4y_2, \quad D_1 := 1 - 4y_2.$$

Let  $F(t) := t_6 f_1(t) + t_7 f_2(t)$  be the Cayley polynomial associated with  $f_1(t)$  and  $f_2(t)$ . Its support polytope is the Cayley polytope  $\tilde{\Delta} = \Delta_1 * \Delta_2 \subset \tilde{M}_{\mathbb{R}} = M_{\mathbb{R}} \oplus \mathbb{R}^2$  which is the convex hull of the vectors:

$$\begin{aligned} u_1 &= (0, 0, 0, 0, 0; 1, 0), \quad u_2 = (1, 0, 0, 0, 0; 1, 0), \quad u_3 = (0, 1, 0, 0, 0; 1, 0), \\ u_4 &= (0, 0, 1, 0, 0; 1, 0), \quad u_5 = (0, 0, 0, 1, 0; 1, 0), \quad u_6 = (0, 0, 0, 0, 0; 0, 1), \\ u_7 &= (-1, -1, -1, -1, 0; 0, 1), \quad u_8 = (0, 0, 0, 0, 1; 0, 1), \quad u_9 = (0, 0, 0, 0, -1; 0, 1). \end{aligned}$$

**Proposition 7.11.** *Let  $A := \{u_1, \dots, u_9\} \subset \tilde{M}$ . Then the principal  $A$ -determinant of  $F(t)$  has the following form:*

$$\begin{aligned} E_A(F) &= a_1^8 a_2^8 a_3^8 a_4^8 a_5^5 a_6^5 a_7^4 D_0 D_1^4 = \\ &-640a_1^8 a_2^8 a_3^8 a_4^8 a_5^8 a_6^8 a_7^8 - 16777216a_1^{10} a_2^{10} a_3^{10} a_4^{10} a_5^{10} a_6^8 a_7^8 + \\ &160a_1^8 a_2^8 a_3^8 a_4^8 a_5^8 a_6^7 a_7^7 + \underline{16777216a_1^{10} a_2^{10} a_3^{10} a_4^{10} a_5^{10} a_6^9 a_7^9} + \\ &\underline{65536a_1^{10} a_2^{10} a_3^{10} a_4^{10} a_5^{10} a_6^5 a_7^5} + 6291456a_1^{10} a_2^{10} a_3^{10} a_4^{10} a_5^{10} a_6^7 a_7^7 - \\ &1048576a_1^{10} a_2^{10} a_3^{10} a_4^{10} a_5^{10} a_6^6 a_7^6 - \underline{1024a_1^8 a_2^8 a_3^8 a_4^8 a_5^8 a_6^{10} a_7^{10}} + \\ &1280a_1^8 a_2^8 a_3^8 a_4^8 a_5^8 a_6^9 a_7^9 - 512a_1^9 a_2^9 a_3^9 a_4^9 a_5^9 a_6^5 a_7^5 - \\ &20a_1^8 a_2^8 a_3^8 a_4^8 a_5^8 a_6^6 a_7^6 + 131072a_1^9 a_2^9 a_3^9 a_4^9 a_5^9 a_6^8 a_7^8 + \\ &\underline{a_1^8 a_2^8 a_3^8 a_4^8 a_5^8 a_6^5 a_7^5} + 8192a_1^9 a_2^9 a_3^9 a_4^9 a_5^9 a_6^6 a_7^6 - \\ &49152a_1^9 a_2^9 a_3^9 a_4^9 a_5^9 a_6^7 a_7^7 - 131072a_1^9 a_2^9 a_3^9 a_4^9 a_5^9 a_6^9 a_7^9, \end{aligned}$$

where the terms corresponding to the vertices of the Newton polytope of  $E_A(F)$  are underlined.

*Proof.* The principal  $A$ -determinant can be found by using the algorithm proposed by A. Dickenstein and B. Sturmfels [DS] via the computation of the corresponding Chow forms.  $\square$

*Remark 7.12.* We note that  $D_0$  is the principal component of the discriminant locus  $E_A(F) = 0$  and the component  $D_1$  corresponds to the edge  $\Gamma$  of  $\tilde{\Delta}$  with

$$\Gamma \cap \tilde{M} = \{u_6, u_8, u_9\} = \{(0, 0, 0, 0, 0; 0, 1), (0, 0, 0, 0, 1; 0, 1), (0, 0, 0, 0, -1; 0, 1)\}.$$

The Newton polytope of  $E_A(F)$  is the secondary polytope  $\text{Sec}(A)$  depicted in Figure 1. The vertices of  $\text{Sec}(A)$  are in one-to-one correspondence with coherent triangulations  $\mathcal{T}_1, \dots, \mathcal{T}_4$  of  $\tilde{\Delta}$  which are:

$$\begin{aligned}
\mathcal{T}_1 &= \{ \langle u_1, u_3, u_4, u_5, u_6, u_7, u_8 \rangle, \langle u_1, u_2, u_4, u_5, u_6, u_7, u_8 \rangle, \langle u_1, u_2, u_3, u_5, u_6, u_7, u_8 \rangle, \\
&\quad \langle u_1, u_2, u_3, u_4, u_6, u_7, u_8 \rangle, \langle u_1, u_2, u_3, u_4, u_5, u_6, u_8 \rangle, \langle u_1, u_3, u_4, u_5, u_6, u_7, u_9 \rangle, \\
&\quad \langle u_1, u_2, u_4, u_5, u_6, u_7, u_9 \rangle, \langle u_1, u_2, u_3, u_5, u_6, u_7, u_9 \rangle, \langle u_1, u_2, u_3, u_4, u_6, u_7, u_9 \rangle, \\
&\quad \langle u_1, u_2, u_3, u_4, u_5, u_6, u_9 \rangle \}. \\
\mathcal{T}_2 &= \{ \langle u_1, u_3, u_4, u_5, u_7, u_8, u_9 \rangle, \langle u_1, u_2, u_4, u_5, u_7, u_8, u_9 \rangle, \langle u_1, u_2, u_3, u_5, u_7, u_8, u_9 \rangle, \\
&\quad \langle u_1, u_2, u_3, u_4, u_7, u_8, u_9 \rangle, \langle u_1, u_2, u_3, u_4, u_5, u_8, u_9 \rangle \}. \\
\mathcal{T}_3 &= \{ \langle u_2, u_3, u_4, u_5, u_7, u_8, u_9 \rangle, \langle u_1, u_2, u_3, u_4, u_5, u_7, u_9 \rangle, \langle u_1, u_2, u_3, u_4, u_5, u_7, u_8 \rangle \}. \\
\mathcal{T}_4 &= \{ \langle u_1, u_2, u_3, u_4, u_5, u_7, u_8 \rangle, \langle u_1, u_2, u_3, u_4, u_5, u_7, u_9 \rangle, \langle u_2, u_3, u_4, u_5, u_6, u_7, u_8 \rangle, \\
&\quad \langle u_2, u_3, u_4, u_5, u_6, u_7, u_9 \rangle \}.
\end{aligned}$$




FIGURE 1. Secondary polytope and coherent triangulations

The generating function  $I_P(y)$  for intersection numbers corresponding to the monomial  $P(x) = x_1^{k_1} x_2^{k_2}$  can be computed from the intersection theory on the Morrison-Plesser moduli spaces. Using the two independent integral relations between  $v_1, \dots, v_7$

$$v_1 + \dots + v_5 = 0, \quad v_6 + v_7 = 0,$$

we see that the Mori cone  $K_{\text{eff}}(\mathbb{P})$  is spanned by two vectors

$$l^{(1)} = (1, 1, 1, 1, 1, 0, 0), \quad l^{(2)} = (0, 0, 0, 0, 0, 1, 1).$$

The Morrison-Plesser moduli spaces are  $\mathbb{P}_\beta = \mathbb{P}^{5b_1+4} \times \mathbb{P}^{2b_2+1}$  ( $b_1, b_2 \in \mathbb{Z}_{\geq 0}$ ). The cohomology of  $\mathbb{P}_\beta$  are generated by the hyperplane classes  $[H_1]$  and  $[H_2]$ . Let  $E_1 := [H_1] + 2[H_2]$  and  $E_2 := 4[H_1]$ . Then the nef-partition  $\Delta = \Delta_1 + \Delta_2$  of polytopes induces the nef-partition of the anticanonical divisor

$$-K_{\mathbb{P}} = E_1 + E_2 = ([H_1] + 2[H_2]) + (4[H_1]).$$

It is straightforward to see that the corresponding Morrison-Plesser class is

$$\Phi_\beta = ([H_1] + 2[H_2])^{b_1+2b_2} (4[H_1])^{4b_1}.$$

So we get

$$I_P(y) = \sum_{b_1, b_2 \geq 0} \langle [H_1]^{k_1} [H_2]^{k_2} ([H_1] + 2[H_2])^{b_1+2b_2+1} (4[H_1])^{4b_1+1} \rangle_\beta y_1^{b_1} y_2^{b_2}.$$

Using the intersection theory on  $\mathbb{P}_\beta$ , we obtain

$$I_P(y) = \sum_{b_1, b_2 \geq 0} 2^{8b_1+2b_2-k_2+3} \frac{(b_1 + 2b_2 + 1)!}{(b_1 - k_1 + 3)!(2b_2 - k_2 + 1)!} y_1^{b_1} y_2^{b_2}.$$

By Theorem 6.1 the function  $I_P(y)$  admits the integral representation:

$$I_P(y) = \frac{1}{(2\pi i)^2} \int_\Gamma \frac{4z_1^{k_1+1} z_2^{k_2} (z_1 + 2z_2) dz_1 \wedge dz_2}{(z_1^5 - (z_1 + 2z_2)(4z_1)^4 y_1)(z_2^2 - (z_1 + 2z_2)^2 y_2)}$$

with the cycle

$$\Gamma = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1^5 - (z_1 + 2z_2)(4z_1)^4 y_1| = |z_2^2 - (z_1 + 2z_2)^2 y_2| = \delta\},$$

where  $\delta$  is positive. These integrals can be easily computed and yield the same rational functions as in Proposition 7.10.

## REFERENCES

- [AY] L.A. Aizenberg, A.P. Yuzhakov, *Integral representations and residues in multidimensional complex analysis*, Trans. of Math. Monographs **58**, AMS 1983.
- [Bat] V.V. Batyrev, *Quantum cohomology rings of toric manifolds*, Asterisque **218** (1993), 9–34.
- [BB1] V.V. Batyrev, L.A. Borisov, *Dual cones and mirror symmetry for generalized Calabi-Yau manifolds*. In Mirror Symmetry II (ed. S.-T. Yau) (1995), 65–80.
- [BB2] V.V. Batyrev, L.A. Borisov, *Mirror duality and string-theoretic Hodge numbers*, Invent. math. **126** (1996), 183–203.
- [BM1] V.V. Batyrev, E.N. Materov, *Toric residues and mirror symmetry*, Moscow Mathematical Journal **2** (2002), No. 3, 435–475.
- [BM2] V.V. Batyrev, E.N. Materov, *Toric residues and Grothendieck residues*, in preparation.
- [BvS] V.V. Batyrev, D. van Straten, *Generalized hypergeometric functions and rational curves on Calabi-Yau complete intersections in toric varieties*, Commun. Math. Phys. **168** (1995), 493–533.
- [Bo] L.A. Borisov, *Towards the mirror symmetry for Calabi-Yau complete intersections in Gorenstein toric Fano varieties*.
- [CDS] E. Cattani, A. Dickenstein and B. Sturmfels, *Residues and resultants*, J. Math. Sci. Univ. Tokyo **5** (1998), 119–148.
- [Cox] D. Cox, *Toric residues*, Ark. Mat. **34** (1996), 73–96.
- [DS] A. Dickenstein, B. Sturmfels, *Elimination theory in codimension two*, J. Symbolic Comput. **34** (2002), 119–135.
- [GKZ] I.M. Gelfand, M.M. Kapranov and A.V. Zelevinsky, *Discriminants, Resultants and Multidimensional Determinants*, Birkhäuser, Boston, 1994.

- [HKTY] S. Hosono, A. Klemm, S. Theisen, S.-T. Yau, *Mirror symmetry, mirror map and applications to complete intersection Calabi-Yau spaces*, Nuclear Phys. B **433** (1995), no. 3, 501–552.
- [DK] V.I. Danilov, A.G. Khovanskii, *Newton polyhedra and an algorithm for computing Hodge-Deligne numbers*, Math. USSR Izvestiya **29** (1987), 279–298.
- [Mav1] A.R. Mavlyutov, *Cohomology of complete intersections in toric varieties*, Pacific J. of Math., vol 191 (1999), no 1, 133–144.
- [Mav2] A.R. Mavlyutov, *Semiample hypersurfaces in toric varieties*, Duke Math. J. **101** (2000), 85–116.
- [MP] D. Morrison, M. Plesser, *Summing the instantons: Quantum cohomology and mirror symmetry in toric varieties*, Nucl. Phys. B **440** (1995), 279–354.
- [Ts] A.K. Tsikh, *Multidimensional Residues and their Applications*, Trans. of Math. Monographs **103**, AMS 1992.

MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE 10, TÜBINGEN  
D-72076, GERMANY

*E-mail address:* victor.batyrev@uni-tuebingen.de

MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE 10, TÜBINGEN  
D-72076, GERMANY

*E-mail address:* evgeny.materov@uni-tuebingen.de